

Environmental Justice Index Indicators

Overall Environmental Justice Rank	Social Vulnerability	Racial/ Ethnic Minority Status	Minority Status
		Socioeconomic Status	Poverty
			No High School Diploma
			Unemployment
			Housing Tenure
			Housing Burdened Lower-Income Households
			Lack of Health Insurance
			Lack of Broadband Access
		Household Characteristics	Age 65 and Older
			Age 17 and Younger
			Civilian with a Disability
			Speaks English "Less than Well"
	Housing Type	Group Quarters	
		Mobile Homes	
	Environmental Burden	Air Pollution	Ozone
			PM2.5
			Diesel Particulate Matter
			Air Toxics Cancer Risk
		Potentially Hazardous & Toxic Sites	National Priority List Sites
			Toxic Release Inventory Sites
			Treatment, Storage, and Disposal Sites
			Risk Management Plan Sites
			Coal Mines
			Lead Mines
Built Environment		Recreational Parks	
		Houses Built Pre-1980	
		Walkability	
Transportation Infrastructure		High-Volume Roads	
		Railways	
		Airports	
Water Pollution	Impaired Surface Water		
Health Vulnerability	Pre-existing Chronic Disease Burden	Asthma*	
		Cancer*	
		High Blood Pressure*	
		Diabetes*	
		Poor Mental Health*	

TEXT-ONLY VERSION

Social vulnerability module

- Racial/Ethnic Minority Status
 - » Minority Status
- Socioeconomic Status
 - » Poverty
 - » No High School Diploma
 - » Unemployment
 - » Housing Tenure
 - » Housing Burdened Lower-Income Households
 - » Lack of Health Insurance
 - » Lack of Broadband Access
- Household Characteristics
 - » Age 65 and Older
 - » Age 17 and Younger
 - » Civilian with a Disability
 - » Speaks English "Less than Well"
- Housing Type
 - » Group Quarters
 - » Mobile Homes

Environmental Burden Module

- Air Pollution
 - » Ozone
 - » PM2.5
 - » Diesel Particulate Matter
 - » Air Toxics Cancer Risk
- Potentially Hazardous and Toxic Sites
 - » National Priority List Sites
 - » Toxic Release Inventory Sites
 - » Treatment, Storage, and Disposal Sites
 - » Risk Management Plan Sites
 - » Coal Mines
 - » Lead Mines
- Built Environment
 - » Recreational Parks
 - » Houses Built Pre-1980
 - » Walkability
- Transportation Infrastructure
 - » High-Volume Roads
 - » Railways
 - » Airports
- Water Pollution
 - » Impaired Surface Water

Health Vulnerability Module

- Pre-existing Chronic Disease Burden
 - » Asthma*
 - » Cancer*
 - » High Blood Pressure*
 - » Diabetes*
 - » Poor Mental Health*

Health vulnerability measures are marked with asterisks because they are calculated differently than other indicators. While most indicators can have a range of values, the health vulnerability indicators represent only whether a given census tract experiences a high estimated prevalence of disease or not. See the technical documentation for more information on indicators and index scoring.



Environmental Burden Module

Cumulative environmental burden can be understood as the sum of activities that cause environmental pollution or negatively affect environmental and human health (Owusu et al. 2022). The approach taken here to quantify cumulative environmental burden includes assessments of both features of the environment that contribute to good health (salutogenic features) and features of the environment that may be detrimental to human health (pathogenic features). While many cumulative impacts and EJ mapping tools consider only pathogenic features of the environment (California Office of Environmental Health Hazard Assessment, 2021; Min et al., 2019; United States Environmental Protection Agency, 2019), a growing body of literature has documented the importance of salutogenic features in determining environmental quality and measuring health disparities attributable to environmental conditions (Brulle & Pellow, 2006; Maizlish et al., 2019; Pastor et al., 2005; Shrestha et al., 2016).

Air Pollution: Ozone

Indicator: Mean annual number of days with maximum 8-hour average ozone concentration over the National Ambient Air Quality Standard (NAAQS), averaged over three years (2014-2016)

Data Year: 2014-2016

Data source: U.S. Environmental Protection Agency Air Quality System (AQS; combined monitoring and modeled data)

Rationale:

Both acute and long-term exposure to elevated levels of ozone in air are associated with negative health effects ranging from increased morbidity and mortality due to respiratory and cardiovascular disease (Crouse et al., 2015; Last et al., 2017). Together with PM_{2.5}, ozone is a major contributor to air pollution-related morbidity and mortality, with an estimated 4,700 ozone-related deaths in the United States in 2005 (Fann et al., 2012).

Air Pollution: PM_{2.5}

Indicator: Mean annual percent of days with daily 24-hour average PM_{2.5} concentrations over the National Ambient Air Quality Standard (NAAQS), averaged over three years (2014-2016)

Data Year: 2014-2016

Data source: U.S. Environmental Protection Agency Air Quality System (AQS; combined monitoring and modeled data)

Rationale:

Inhalation of particulate matter with a diameter of 2.5 microns or less (PM_{2.5}) can have a number of adverse effects on health and well-being. Acute exposure to elevated levels of PM_{2.5} can lead to irritation of eyes, nose, throat and lungs, and increases relative risk of acute cardiovascular events including admission to a hospital for stroke (Rajagopalan et al., 2018). Long-term exposure to elevated levels of PM_{2.5} is associated with higher rates of mortality from a number of conditions ranging from cancer to cardiopulmonary disease (Dockery & Pope, 1994). In the U.S. in 2005, an estimated 130,000 deaths were attributable to PM_{2.5}-related causes (Fann et al., 2012).

Air Pollution: Diesel Particulate Matter

Indicator: Diesel particulate matter concentrations in air, $\mu\text{g}/\text{m}^3$

Data Year: 2014

Data source: U.S. Environmental Protection Agency National Air Toxics Assessment (NATA: modeled data)

Rationale:

Diesel particulate matter (DPM) is a particle emission from a diesel motor made of an elemental carbon core and various adsorbed organics compounds and other chemical components (Wichmann, 2007). Evidence indicates that DPM exposure may cause respiratory symptoms via inflammation and oxidative stress (Ristovski et al., 2012). Acute exposure to DPM has been associated with acute coronary syndrome (ACS) and other cardiovascular issues (Peters et al., 2001) and DPM contains carcinogens such as benzene and formaldehyde that may lead to the development of certain kinds of cancer (Krivoshto et al., 2008).

Air Pollution: Air Toxics Cancer Risk

Indicator: Lifetime cancer risk from inhalation of air toxics

Data Year: 2014

Data source: U.S. Environmental Protection Agency National Air Toxics Assessment (NATA; modeled data)

Rationale:

Air toxics cancer risk is a composite measure assessing the cancer risk associated with inhaling 140 different hazardous air pollutants (HAPs). HAPs such as benzene, dioxin, formaldehyde, and ethylene oxide are known carcinogens which, at various concentrations, contribute to lifetime risk of developing certain types of cancer (Loh et al., 2007; Reynolds et al., 2003; Whitworth et al., 2008; Wu et al., 2009). The cancer risks estimated by NATA are based on modeled exposure concentrations, assessments of each pollutant's unit risk estimate, and inhalation reference concentration. It is important to note that diesel particulate matter (DPM), which is another CDC/ATSDR EJI indicator, is one of the HAPs included in the 2014 NATA lifetime cancer risk model. However, the DPM indicator is represented as distinct from the air toxics cancer risk indicator because DPM is only one of the 140 HAPs used to create the 2014 NATA lifetime cancer risk estimate and is associated with many health issues other than cancer. For more information on the 2014 NATA, including a full list of HAPs included in the lifetime cancer risk model, please visit <https://www.epa.gov/national-air-toxics-assessment/2014-nata-assessment-results>.

Potentially Hazardous & Toxic Sites: National Priority List Sites

Indicator: Proportion of tract area within 1-mi buffer of EPA National Priority List (NPL) sites

Data Year: 2021

Data source: U.S. Environmental Protection Agency Facility Registry Service (FRS)

Rationale:

Sites on the EPA's National Priorities List (NPL), which are designated by the U.S. EPA as priorities through hazard assessment, nomination by states or territories, or issuance of a health advisory by the Agency for Toxic Substance and Disease Registry, can present several potential hazards to the health and well-being of neighboring communities. While actual risks to health vary by sites, proximity to these sites can have important and complex effects on community stress and perceptions of risk (Kiel & Zabel, 2001; Pearsall, 2010). Furthermore, legacy contaminants associated with many of these sites can affect multiple environmental media, becoming airborne with windblown dust or leaching into soil and groundwater and possibly exposing surrounding communities through drinking water or vapor intrusion.

Potentially Hazardous & Toxic Sites: Toxic Release Inventory Sites

Indicator: Proportion of tract area within 1-mi buffer of Toxic Release Inventory (TRI) sites

Data Year: 2021

Data source: U.S. Environmental Protection Agency Facility Registry Service (FRS)

Rationale:

Sites listed through the EPA's Toxic Release Inventory (TRI) include all facilities with 10 or more full time employees which operate within certain industrial sectors and annually either 1) manufacture more than 25,000 pounds of listed chemicals or 2) used more than 10,000 pounds of listed chemicals. These sites can affect the health of neighboring communities through routine chemical releases into air, soil, or water. Residential proximity to TRI sites has been linked to higher rates of hospitalization for COPD (Brown-Amilian & Akolade, 2021) as well as increased risks for certain kinds of cancer (Bulka et al., 2016). Additionally, TRI sites and other noxious and unwanted land uses can produce noise and odor pollution and, particularly in communities burdened by multiple such land uses, can lead to increased burden of community stress (Wilson et al., 2012).

Potentially Hazardous & Toxic Sites: Treatment, Storage, and Disposal Facilities

Indicator: Proportion of tract area within 1-mi buffer of EPA Treatment, Storage, and Disposal Facilities (TSDF)

Data Year: 2021

Data source: U.S. Environmental Protection Agency Facility Registry Service (FRS)

Rationale:

Sites listed as Treatment, Storage, and Disposal Facilities (TSDF) are responsible for handling hazardous wastes such as manufacturing by-products, cleaning fluids, or pesticides throughout the process of collection, transfer, and ultimately disposal. Volatile substances generated by waste may become aerosolized or migrate into soil and water, leading to vapor intrusion or contamination of groundwater (Johnston & MacDonald Gibson, 2015; Marshall et al., 1993). Proximity to hazardous waste sites has been linked to increased rates of hospitalizations for diseases such as stroke, diabetes, and coronary heart disease (Kouznetsova et al., 2007; Sergeev & Carpenter, 2005; Shcherbatykh et al., 2005).

Potentially Hazardous & Toxic Sites: Risk Management Plan Sites

Indicator: Proportion of tract area within 1-mi buffer of EPA Risk Management Plan (RMP) sites

Data Year: 2021

Data source: U.S. Environmental Protection Agency Facility Registry Service (FRS)

Rationale:

The EPA's Risk Management Plan (RMP) program covers ~12,500 of the nation's most high-risk facilities that produce, use, or store significant amounts of certain highly toxic or flammable chemicals. These facilities must prepare plans for responding to a worst-case incident such as a major fire or explosion that releases a toxic chemical into the surrounding community (US Environmental Protection Agency, 2016). There are many negative health effects associated with residing in proximity to RMP sites. The EPA estimates that about 150 "reportable" incidents of unplanned chemical releases occur each year at RMP facilities, separate from the daily toxic emissions that are allowed under most operating permits. The EPA notes that these incidents "pose a risk to neighboring communities and workers because they result in fatalities, injuries, significant property damage, evacuations, sheltering in place, or environmental damage" (US Environmental

Protection Agency, 2021). Besides direct deaths and injuries caused by chemical release and explosion incidents, research shows increased risk of cancer and respiratory illness from toxic air pollution exposure at these sites. Although the effects of proximity to RMP sites on community stress has not formally been assessed, it is also reasonable to assume that fear of potential chemical plant disasters contributes to the burden of psychosocial stress imposed on communities by cumulative environmental and social stressors (Hynes & Lopez, 2007).

Potentially Hazardous & Toxic Sites: Coal Mines

Indicator: Proportion of tract area within 1-mi buffer of a coal mine

Data Year: 2021

Data source: U.S. Mine Safety and Health Administration Mine Data Retrieval System (MDRS)

Rationale:

Coal mining, while on the decline in the United States, is still of substantial concern for the health of exposed communities, including both traditional underground mining methods and surface mining methods, such as mountaintop removal (MTR). Studies have observed elevated blood inflammation levels, increased cardiopulmonary, lung, and kidney disease, and increased rates of lung cancer mortality in heavy Appalachian coal mining communities as a result of air pollution from mining activity (Hendryx et al., 2010; Hendryx & Ahern, 2008; Hendryx & Luo, 2015). Proximity to MTR sites has been linked to impaired respiratory health, including increased occurrence of chronic obstructive pulmonary disease (COPD)(Hendryx & Luo, 2015) and may predict increased risk for depressive and substance use disorders (Canu et al., 2017). Air pollution from coal mining has also been connected to adverse effects in-utero for pregnant women, including low-birthweight (Ahern et al., 2011). Exposure pathways to coal contamination are also multifactorial. Coal slurry (the practice of disposing liquified coal wastes underground) can leach coal-related pollutants into well and ground water, potential drinking water sources for residents (Ducatman et al., 2010).

Potentially Hazardous & Toxic Sites: Lead Mines

Indicator: Proportion of tract area within 1-mi buffer of an active lead mine

Data Year: 2021

Data source: U.S. Mine Safety and Health Administration Mine Data Retrieval System (MDRS)

Rationale:

Lead mines constitute an important health risk for surrounding communities. Studies in the U.S. have suggested that soil and dust contaminated from lead mining as well as other waste-byproducts of mining pose a health hazard to nearby communities, particularly to children (Malcoe et al., 2002; Murgueytio et al., 1998). Studies outside of the U.S. evaluating health risks associated with communities in close proximity to active lead mines have found evidence of elevated blood lead levels in children (Schirnding et al., 2003; Zhang et al., 2012).

Built Environment: Lack of Recreational Parks

Indicator: Proportion of tract area not within 1-mi buffer of a park, recreational area, or public forest

Data Year: 2020

Data source: TomTom MultiNet® Enterprise Dataset

Rationale:

Parks and greenspaces represent important healthy features of the built environment, providing spaces for physical recreation and promoting physical activity (Bedimo-Rung et al., 2005; Cohen et al., 2007), though evidence that parks promote physical activity in rural areas is mixed (Reuben et al., 2020; Roemmich et al., 2018). Parks and greenspaces also play an important role in mitigating urban heat island effects (P. Lin et al., 2017; Shishegar, 2014) and can offer refuge on extreme heat days (Brown et al., 2015; Voelkel et al., 2018). Proximity and access to parks and greenspaces can also have important implications for mental health, with studies indicating that measures of proximity and access to these spaces are associated with better overall mental health (Bojorquez & Ojeda-Revah, 2018; Sturm & Cohen, 2014; Wood et al., 2017). While park design quality, and neighborhood perceptions of safety can have important mediating effects on these benefits (Cohen et al., 2010; Cutts et al., 2009; Rigolon et al., 2018), and while there are concerns associated with “greening” and gentrification (Mullenbach & Baker, 2020; Wolch et al., 2014), these spaces nearly always provide an overall benefit to neighboring communities and lack of access constitutes an important issue for health and environmental justice (Boone et al., 2009; Jennings et al., 2012; Rigolon, 2017; Rigolon et al., 2018).

Built Environment: Housing Built Pre-1980

Indicator: Proportion of occupied housing units built prior to 1980

Data Year: 2015-2019

Data source: U.S. Census Bureau American Community Survey (ACS)

Rationale:

Age of housing units has important implications for potential exposure to lead. While lead-based paint was banned in 1978, housing built around that time or prior often contain underlying layers of lead-based paint. While underlying layers of lead-based paint do not necessarily constitute a health risk, chipping or flaking that exposes underlying layers of lead-based paints may lead to ingestion by children (Lanphear et al., 1996). Measures of housing built prior to the ban on lead-based paint have repeatedly been identified as one of the leading predictors of blood-lead levels in children (Kim et al., 2002; Sadler et al., 2017; Schultz et al., 2017). There are no known safe levels of lead exposure, especially among children, who are highly susceptible to neurological and developmental issues associated with lead exposure.

Built Environment: Lack of Walkability

Indicator: National Walkability Index Score

Data Year: 2021

Data source: U.S. Environmental Protection Agency National Walkability Index

Rationale:

The U.S. Centers for Disease Control and Prevention (CDC) considers walkability as ‘the idea of quantifying the safety and desirability of the walking routes’ (Smith, 2015). This conceptualization of walkability, stemming from the scientific

evidence that walking can boost metabolism, lower blood sugar and improve mental health (Barton et al., 2009), has become a quantifiable variable to study health-promoting effects of the built environment. Research shows that nearby available locations for walking and biking promote physical activity. Higher residential neighborhood walkability has been associated with more walking, higher overall physical activity, lower body mass index (BMI), lower incidence of diabetes, improved glycemic control among residents, and lower premature mortality (Awuor & Melles, 2019; Chen & Kwan, 2015; L. Frank et al., 2010; L. D. Frank et al., 2004, 2005, 2006; Freeman et al., 2013; Hirsch et al., 2014; US Environmental Protection Agency, 2014, 2017). Measures of neighborhood walkability that include measures of street connectivity, transit stop density, and land use mix, all features of the EPA's National Walkability Index, have also been shown to be positively associated with various measures of accessibility for older adults and persons with disabilities (King et al., 2011; Kwon & Akar, 2022; Mahmood et al., 2020). While it is important to note that the associations between built environment measures of walkability on health may be different in rural and urban neighborhoods (Stowe et al., 2019), and while these measures may not account for physical or social factors that could mediate the effects of walkability on physical activity and health benefits (Bracy et al., 2014; Forsyth, 2015), walkability nevertheless constitutes an important environmental amenity.

Transportation Infrastructure: High Volume Roads

Indicator: Proportion of tract area within 1-mi buffer of a high-volume street or road

Data Year: 2020

Data source: TomTom MultiNet® Enterprise Dataset

Rationale:

High-volume roads, such as interstate highways, can constitute major hazards to surrounding communities. Vehicular emissions are a major source of air pollutants such as ozone and diesel particulate matter, and proximity to busy roads has been associated with a number of adverse respiratory symptoms, childhood cancers, adverse birth outcomes, and overall mortality (Boothe & Shendell, 2008). Water runoff from roads can also lead to deposition of heavy metals and other pollutants in nearby soils and waters (Khalid et al., 2018; Sutherland & Tolosa, 2001). Noise pollution associated with traffic is also associated with significant increases in community stress (Barbaresco et al., 2019) and can lead to elevated risk of cardiovascular disease (Münzel et al., 2021) and adverse mental health outcomes (Díaz et al., 2020).

Transportation Infrastructure: Railways

Indicator details: Proportion of tract area within 1-mi buffer of a railway

Data Year: 2020

Data source: TomTom MultiNet® Enterprise Dataset

Rationale:

Like roads, railways can also present a significant source of noise pollution to nearby communities. This noise pollution can constitute a major annoyance and source of community stress, especially when combined with noise pollution from traffic (Öhrström et al., 2007). Among all transportation-associated sources of noise pollution, railway noise is associated with the most significant levels of sleep disruption and associated increases in stress and diastolic blood pressure (Elmenhorst et al., 2019; Petri et al., 2021).

Transportation Infrastructure: Airports

Indicator: Proportion of tract area within 1-mi buffer of an airport

Data Year: 2020

Data source: TomTom MultiNet® Enterprise Dataset

Rationale:

Airports are important sources of noise pollution. Studies indicate that noise pollution associated with residential proximity to airports can constitute a significant nuisance, and can lead to elevated levels of stress and sleep disturbance (Elmenhorst et al., 2019; Ogneva-Himmelberger & Cooperman, 2010; Ozkurt et al., 2015). Airports are also important sources of air, soil, and groundwater contamination. Accidental releases from leaky storage tanks, use of hazardous chemicals in rescue and firefighting training, and stormwater runoff all contribute to infiltration of chemicals such as benzene, trichloroethylene, carbon tetrachloride, and a range of perfluorochemicals into soil and groundwater (Nunes et al., 2011).

Water Pollution: Impaired Surface Water

Indicator: Percent of tract watershed area classified as impaired

Data Year: 2019

Data source: U.S. Environmental Protection Agency Watershed Index Online (WSIO)

Rationale:

Surface waters such as rivers and lakes are important for recreation and fishing, and impairment of these waters can constitute a potential nuisance or even hazard to nearby residents. Waters may be classified as impaired due to elevated levels of waterborne pathogens or significant contamination by toxic substances. Waterborne pathogens can pose a significant health risk through recreational exposure (McKee & Cruz, 2021), while ingestion of fish from chemically-impaired waters can be a significant exposure pathway for a number of pollutants that bioaccumulate in tissues (Dórea, 2008).

Social Vulnerability Module

Literature regarding environmental injustice documents the disproportionate placement of hazardous waste sites, industrial facilities, busy roads and railways, and sewage treatment plants in socially vulnerable neighborhoods (Bullard et al., 2008; Mohai et al., 2009; Mohai & Saha, 2007; Morello-Frosch et al., 2011). These communities are thus more likely to be exposed to harmful pollutants and experience poor health outcomes, such as cardiovascular disease, asthma, perinatal outcomes, and mental health impacts (Morello-Frosch et al., 2011). Given the inequities associated with social vulnerability, these communities are also less likely to receive financial assistance for environmental and disaster recovery, have access to mental and physical health services (Tate & Emrich, 2021), or have the social capital or resources to influence environmental decision-making (Pearsall, 2010). Thus, socially vulnerable communities may be particularly vulnerable to procedural environmental injustices.

Racial/Ethnic Minority Status: Minority Status

Indicator: Percent of population that is a racial/ethnic minority (all persons except white, non-Hispanic)

Data Year: 2015-2019

Data source: U.S. Census Bureau American Community Survey (ACS)

Rationale:

Historical and ongoing racial residential segregation, race-related income inequality, and other forms of institutional and systemic racism have often limited the ability of these populations to advocate against unwanted land uses or influence environmental decision-making, as borne out by the disproportionate location of contamination sites near non-white populations (Bullard et al., 2008; Cutter et al., 2003; Ernst, 1994; Lee, 1992). Systemic racism has been labeled by the CDC as a serious public health threat (see statement here: <https://www.cdc.gov/media/releases/2021/s0408-racism-health.html>). A growing body of data suggest that aspects of systemic and structural racism contribute to health disparities, including those associated with environmental pollution, through a number of pathways, including discrimination by the institutional medical system (Boateng & Aslakson, 2021). Minorities experiencing negative health effects associated with environmental pollution may experience barriers to accessing health care due to discrimination and other factors and may suffer disproportionately adverse outcomes (Neighbors et al., 2007; Smedley, 2012; Williams & Mohammed, 2009).

Socioeconomic Status: Poverty

Indicator: Percent of population with income below 200% of federal poverty level

Data Year: 2015-2019

Data source: U.S. Census Bureau American Community Survey (ACS)

Rationale:

Poverty is an indication of economic hardship. Lack of financial resources may hinder a community's ability to influence environmental decision-making, leading to contamination sites being disproportionately located in impoverished areas (Mohai & Bryant, 1991; Mohai & Saha, 2015; Tanzer et al., 2019). Low-income populations are also particularly susceptible to adverse health outcomes, at least in part due to psychosocial and chronic stress and lack of healthcare access (Evans & Kim, 2013; Haushofer & Fehr, 2014; Wright et al., 1998). Research indicates that negative effects of air pollution on birth outcomes are greater for mothers from low-income neighborhoods (Padula et al., 2014; Yi et al., 2010).

Socioeconomic Status: No High School Diploma

Indicator: Percent of population (age 25+) with no high school diploma

Data Year: 2015-2019

Data source: U.S. Census Bureau American Community Survey (ACS)

Rationale:

Educational attainment is an important factor of socioeconomic status and may influence communities' ability to navigate information about pollution, environmental law, and community-scale resources to influence environmental decision-making (Helfand & Peyton, 1999). Education also influences populations' susceptibility to health impacts of negative environmental conditions. Low educational attainment has been shown to be associated with increased risk of adverse birth outcomes (Gray et al., 2014; Thayamballi et al., 2021) and overall mortality (Kan et al., 2008).

Socioeconomic Status: Unemployment

Indicator: Percent of population age 16 and older who are unemployed

Data Year: 2015-2019

Data source: U.S. Census Bureau American Community Survey (ACS)

Rationale:

Unemployment is an important marker of community socioeconomic status. Lack of employment often means limited financial resources as well as decreased social capital due to stigma. These factors can reduce this population's ability to influence environmental decision-making. Furthermore, fear of unemployment can prevent communities from advocating against unwanted land uses that provide employment opportunities (Bullard, 1993), and communities with high rates of unemployment may be more receptive to incoming industrial facilities that offer jobs, essentially trading employment for environmental pollution to avoid extreme poverty (Shrader-Frechette, 2002). Unemployment is also associated with stress and stress-related inflammation, potentially rendering these populations more vulnerable to health effects mediated by stress (Ala-Mursula et al., 2013; Dettenborn et al., 2010; Heikkala et al., 2020).

Socioeconomic Status: Housing Tenure

Indicator: Percent of housing units that are renter-occupied

Data Year: 2015-2019

Data source: U.S. Census Bureau American Community Survey (ACS)

Rationale:

Renters are often seen as more transitory and, thus, may have less social capital within the context of environmental decision-making, especially within the context of environmental efforts specifically geared towards homeowners who have vested rights and interests in defending local environmental quality and land values (Perkins et al., 2004; Shapiro, 2005). Additionally, research consistently supports the idea that renters experience worse health outcomes associated with a range of conditions when compared to homeowners, likely due to complex interactions between general socioeconomic status associated with housing tenure and aspects of the physical and meaning-based environments represented by rented and owned housing units (Hiscock et al., 2003; Mawhorter et al., 2021).

Socioeconomic Status: Housing Burdened, Lower-Income Households

Indicator: Percent of households with annual income less than \$75,000 who are considered burdened by housing costs (pay greater than 30% of monthly income on housing expenses)

Data Year: 2015-2019

Data source: U.S. Census Bureau American Community Survey (ACS)

Rationale:

The U.S. Department of Housing and Urban Development (HUD) and the U.S. Census Bureau define a household as "housing cost burdened" if that household pays greater than 30% of monthly income on housing costs. Housing costs represent a significant financial burden for most households, and populations burdened by housing costs and associated debt may lack financial resources or time to devote to improving environmental conditions. Additionally, research indicates that persons experiencing housing burden may be less likely to have access to preventative care or to postpone health care (Meltzer & Schwartz, 2016). Instability associated with housing cost burden can also exacerbate issues of stress and poor mental health and are correlated with worse developmental and educational outcomes for children (Burgard et al., 2012; Newman & Holupka, 2016; Suglia et al., 2011).

Socioeconomic Status: Lack of Health Insurance

Indicator: Percent of civilian, noninstitutionalized population who have no health insurance

Data Year: 2015-2019

Data source: U.S. Census Bureau American Community Survey (ACS)

Rationale:

The total population of insured persons in the US has consistently declined since 1997, despite the recent uptick in 2018-2019, where about 11% of the population at the time remained uninsured. This population of uninsured persons are commonly of families with low income (with typically one person working in the family), people of color, and undocumented immigrants (Tolbert et al., 2020). Financial burdens associated with healthcare may the reduce uninsured populations' ability to engage in the environmental decision-making process. Further, individuals without insurance have barriers to accessing preventative care following adverse environmental events, increasing risk of morbidity and mortality among uninsured populations (Mulchandani et al., 2019; Woolhandler & Himmelstein, 2017).

Socioeconomic Status: Lack of Internet Access

Indicator: Percent of households with no internet subscription

Data Year: 2015-2019

Data source: U.S. Census Bureau American Community Survey (ACS)

Rationale:

Lack of access to broadband services can impede populations' ability to be engaged in decision-making and to be informed on environmental issues in their communities. The inability to access the internet can also be an important communication barrier during environmental emergencies, for which outreach through internet sources can be a key strategy for public health officials (Houston et al., 2015; Jha et al., 2016; Nguyen et al., 2017; Wong et al., 2017).

Household Characteristics: Age 65 and Older

Indicator: Percent of population aged 65 and older

Data Year: 2015-2019

Data source: U.S. Census Bureau American Community Survey (ACS)

Rationale:

Adults aged 65 and older face higher rates of social isolation than the general population, which can affect their ability to affect change or influence environmental decision-making in their communities (Andrew & Keefe, 2014). Additionally, older populations may be more susceptible to environmental pollution due to lowered immune function and accumulated oxidative stress associated with a lifetime of exposures (Cakmak et al., 2007; Hong, 2013; Morello-Frosch et al., 2011).

Household Characteristics: Age 17 and Younger

Indicator: Percent of population aged 17 and younger

Data Year: 2015-2019

Data source: U.S. Census Bureau American Community Survey (ACS)

Rationale:

Persons below voting age have a limited ability to influence environmental decision-making as well as limited resources, knowledge, or life experiences necessary to affect change (Flanagan et al., 2011). Additionally, children are particularly susceptible to negative health effects associated with a range of environmental pollution due to a combination of physiological sensitivity and behaviors that put them at greater risk (Morello-Frosch et al., 2011). Physiological factors, such as rates of absorption, distribution, metabolism, and excretion of chemicals, make children more vulnerable to environmental pollution than adults (Faustman et al., 2000).

Household Characteristics: Civilian with a Disability

Indicator: Percent of civilian, noninstitutionalized population with a disability

Data Year: 2015-2019

Data source: U.S. Census Bureau American Community Survey (ACS)

Rationale:

Those living with a disability may experience social or physiological barriers to full participation in the environmental decision-making process. Persons with disabilities are often disproportionately affected at every stage of disaster events and disaster recovery (Chakraborty et al., 2019; Peek & Stough, 2010). Furthermore, certain types of disability are associated with increased physiological susceptibility to environmental pollution, particularly PM2.5 and other forms of air pollution (Dales & Cakmak, 2016; H. Lin et al., 2017; Weuve et al., 2016).

Household Characteristics: Speaks English “Less than Well”

Indicator: Percent of persons (age 5 and older) who speak English “less than well”

Data Year: 2015-2019

Data source: U.S. Census Bureau American Community Survey (ACS)

Rationale:

The ability to communicate in English can be an important factor in determining a community's ability to participate in civil discourse surrounding environmental decision-making. Documents and news sources covering environmental issues are often not available in languages other than English, hampering non-English speakers' ability to inform themselves and engage in these issues (Teron, 2016). Furthermore, discrimination against non-English speakers can lead to exclusion from decision making and is correlated with increased stress and reduced quality of life (Gee & Ponce, 2010). Non-English speakers may also be more vulnerable during disasters or extreme climate events if materials aimed at dissemination of emergency information are available only in English (Nepal et al., 2012; White-Newsome et al., 2009).

Housing Type: Group Quarters

Indicator: Percentage of persons living in group quarters (includes college residence halls, residential treatment centers, group homes, military barracks, correctional facilities, and worker's dormitories)

Data Year: 2015-2019

Data source: U.S. Census Bureau American Community Survey (ACS)

Rationale:

Institutionalized persons, those in correctional facilities, nursing homes, and mental hospitals, are particularly vulnerable to environmental injustice and often have limited ability to influence environmental decision-making. For example, persons who are incarcerated or detained often face disproportionate exposures to environmental contaminants due to poor institutional conditions, exposures through hazardous work programs, and a lack of social capital to improve conditions for themselves (Pellow, 2021). Persons institutionalized in nursing homes or mental hospitals face similar issues of autonomy and lack of social capital or physical ability to influence environmental decision-making. Furthermore, persons in institutional facilities are often neglected in environmental decision making and hazard response (Cutter, 2012).

Non-institutionalized persons living in group quarters are also vulnerable to environmental injustice, though perhaps not as clearly as institutionalized persons. Military bases share some characteristics with correctional facilities in that they are often sites of concentrated environmental contamination and many of their residents come from similar socioeconomic backgrounds and have similarly little influence over the day to day operations that result in contamination (Broomandi et al., 2020). People living in group homes, missions, and shelters may have limited legal status, limited time, and limited resources, and thus diminished ability to influence environmental decision-making (Goodling, 2020).

Housing Type: Mobile Homes

Indicator: Percentage of total housing units designated as mobile homes

Data Year: 2015-2019

Data source: U.S. Census Bureau American Community Survey (ACS)

Rationale:

Mobile homes are often clustered in communities confined to low-value areas due to zoning laws and stigma (Maantay, 2002). Mobile homes are also often inhabited by farm workers, who are beholden to landowners for environmental decision-making such as use of agricultural pesticides (Early et al., 2006). These aspects of stigma, zoning, and lack of land ownership can inhibit these populations' ability to influence local environmental policy. Furthermore, issues of poor construction and energy inefficiency can render residents of mobile homes more susceptible to negative health effects associated with air pollution (MacTavish et al., 2006) and extreme heat (Phillips et al., 2021), while observed unreliability of access to drinking water poses further risks to residents' health (Pierce & Jimenez, 2015).

Health Vulnerability Module

High Estimated Prevalence of Asthma

Indicator: Estimated prevalence of asthma among adults 18 and older greater than for 66.66% of U.S. census tracts (2020)

Data Year: 2020

Data source: U.S. Centers for Disease Control and Prevention PLACES Estimates

Rationale:

Outdoor air pollution is associated with increases in asthma attacks (Centers for Disease Control and Prevention, 2020; Peel et al., 2005) and asthma-related ED visits (Norris et al., 1999; Slaughter et al., 2005; Stieb et al., 1996; P. E. Tolbert et al., 2000; Villeneuve et al., 2007). Inhalation of pollutants such as PM2.5, ozone, and diesel particulate matter can lead to oxidative stress which inflames the airways and exacerbates asthma symptoms, and both acute and long-term exposure to asthma are associated with worsening asthma symptoms (Guarnieri & Balmes, 2014).

High Estimated Prevalence of Cancer

Indicator: Estimated prevalence of all-cause cancer (excluding skin cancer) among adults 18 and older greater than for 66.66% of U.S. census tracts (2020)

Data Year: 2020

Data source: U.S. Centers for Disease Control and Prevention PLACES Estimates

Rationale:

Increases in PM2.5 are also associated with increased all-cause mortality for young adult cancer patients diagnosed with all cancer types (Ou et al., 2020). Long-term exposure to PM2.5, ozone, and other air pollutants is associated with increased morbidity and mortality in persons diagnosed with cancer, including lung cancer (Jerrett et al., 2013; Pope III et al., 2002), liver cancer (Deng et al., 2017), pediatric lymphomas, and CNS tumors (Ou et al., 2020). Experimental research suggests that intermediate to long-term exposure to both fine and coarse particulate matter may accelerate oncogenesis (the formation of tumors) and cause increased expression of inflammation and oncogenesis-related genes in rat brains (Ljubimova et al., 2018).

High Estimated Prevalence of High Blood Pressure

Indicator: Estimated prevalence of high blood pressure among adults ≥ 18 greater than for 66.66% of U.S. census tracts (2020)

Data Year: 2020

Data source: U.S. Centers for Disease Control and Prevention PLACES Estimates

Rationale:

Elevated levels of ambient PM2.5, ozone, and other air pollutants are associated with the increased prevalence and elevated risk of adverse health outcomes like heart attack and overall increases in blood pressure, including hypertension (Coogan et al., 2012; Giorgini et al., 2016; Lee, 2020). Long-term exposure to particulate matter, other traffic-related air pollution, and traffic noise pollution have been associated with increased blood pressure and a higher risk of developing hypertension (Dong et al., 2013; Foraster et al., 2014; Fuks et al., 2014). Hypertension is an established risk factor for a number of negative cardiovascular health outcomes, including coronary heart disease and stroke, but cardiovascular complications related to high blood pressure can occur before the onset of established hypertension (Go et al., 2013).

High Estimated Prevalence of Diabetes

Indicator: Estimated prevalence of diabetes among adults 18 and older greater than for 66.66% of U.S. census tracts (2020)

Data Year: 2020

Data source: U.S. Centers for Disease Control and Prevention PLACES Estimates

Rationale:

Research suggests that air pollution, such as PM2.5, can cause oxidative stress and inflammation, leading to impairments in insulin signaling associated with diabetes (Meo et al., 2015). PM2.5 is also associated with markers of systemic inflammation in individuals with diabetes (Dubowsky et al., 2006), which may lead to greater risk of diabetes-related negative health outcomes. Proximity to hazardous sites and land use have also been associated with increased risk of hospitalization among individuals with diabetes (Kouznetsova et al., 2007).

High Estimated Prevalence of Poor Mental Health

Indicator: Estimated prevalence of poor mental health for ≥ 14 days among adults 18 and older greater than for 66.66% of U.S. census tracts (2020)

Time Period: 2020

Data source: U.S. Centers for Disease Control and Prevention PLACES Estimates

Rationale:

Poor mental health can be both caused by and exacerbated by negative environmental quality. One study found that residential proximity to industrial activity negatively impacts mental health directly and by mediating individual's perceptions of neighborhood disorder and personal powerlessness, with these effects being most prominent in racial/ethnic minority populations and populations in poverty (Downey & Van Willigen, 2005). Another exploratory study in the U.S. found a strong positive link between exposure to environmental pollution and an increase of prevalence in psychiatric disorders in affected patients (Khan et al., 2019). Poor environmental quality may also affect the quality of life (i.e. the expectation and concern for one's own health and life) negatively through the mediating effects of increased stress and poor sleep (Chang et al., 2020).

References

- Ahern, M., Mullett, M., MacKay, K., & Hamilton, C. (2011). Residence in Coal-Mining Areas and Low-Birth-Weight Outcomes. *Maternal and Child Health Journal*, 15(7), 974–979. <https://doi.org/10.1007/s10995-009-0555-1>
- Ala-Mursula, L., Buxton, J. L., Ek, E., Koiranen, M., Taanila, A., Blakemore, A. I. F., & Järvelin, M.-R. (2013). Long-Term Unemployment Is Associated with Short Telomeres in 31-Year-Old Men: An Observational Study in the Northern Finland Birth Cohort 1966. *PLOS ONE*, 8(11), e80094. <https://doi.org/10.1371/journal.pone.0080094>
- Andrew, M. K., & Keefe, J. M. (2014). Social vulnerability from a social ecology perspective: A cohort study of older adults from the National Population Health Survey of Canada. *BMC Geriatrics*, 14(1), 90. <https://doi.org/10.1186/1471-2318-14-90>
- Awuor, L., & Melles, S. (2019). The influence of environmental and health indicators on premature mortality: An empirical analysis of the City of Toronto's 140 neighborhoods. *Health & Place*, 58, 102155. <https://doi.org/10.1016/j.healthplace.2019.102155>

- Barbaresco, G. Q., Reis, A. V. P., Lopes, G. D. R., Boaventura, L. P., Castro, A. F., Vilanova, T. C. F., Da Cunha Júnior, E. C., Pires, K. C., Pôrto Filho, R., & Pereira, B. B. (2019). Effects of environmental noise pollution on perceived stress and cortisol levels in street vendors. *Journal of Toxicology and Environmental Health, Part A*, 82(5), 331–337. <https://doi.org/10.1080/15287394.2019.1595239>
- Barton, J., Hine, R., & Pretty, J. (2009). The health benefits of walking in greenspaces of high natural and heritage value. *Journal of Integrative Environmental Sciences*, 6(4), 261–278. <https://doi.org/10.1080/19438150903378425>
- Bedimo-Rung, A. L., Mowen, A. J., & Cohen, D. A. (2005). The significance of parks to physical activity and public health: A conceptual model. *American Journal of Preventive Medicine*, 28(2, Supplement 2), 159–168. <https://doi.org/10.1016/j.amepre.2004.10.024>
- Boateng, A., & Aslakson, R. A. (2021). Navigating the Complex Ecosystem of Race, Ethnicity, Structural Racism, Socioeconomic Factors, Medical Care Delivery, and End-of-Life Care—Casting Away the Compass to Make a Map. *JAMA Network Open*, 4(9), e2126348. <https://doi.org/10.1001/jamanetworkopen.2021.26348>
- Bojorquez, I., & Ojeda-Revah, L. (2018). Urban public parks and mental health in adult women: Mediating and moderating factors. *International Journal of Social Psychiatry*, 64(7), 637–646. <https://doi.org/10.1177/0020764018795198>
- Boone, C. G., Buckley, G. L., Grove, J. M., & Sister, C. (2009). Parks and People: An Environmental Justice Inquiry in Baltimore, Maryland. *Annals of the Association of American Geographers*, 99(4), 767–787. <https://doi.org/10.1080/00045600903102949>
- Boothe, V. L., & Shendell, D. G. (2008). Potential Health Effects Associated with Residential Proximity to Freeways and Primary Roads: Review of Scientific Literature, 1999–2006. *Journal of Environmental Health*, 70(8), 33–41.
- Bracy, N. L., Millstein, R. A., Carlson, J. A., Conway, T. L., Sallis, J. F., Saelens, B. E., Kerr, J., Cain, K. L., Frank, L. D., & King, A. C. (2014). Is the relationship between the built environment and physical activity moderated by perceptions of crime and safety? *International Journal of Behavioral Nutrition and Physical Activity*, 11(1), 24. <https://doi.org/10.1186/1479-5868-11-24>
- Broomandi, P., Guney, M., Kim, J. R., & Karaca, F. (2020). Soil Contamination in Areas Impacted by Military Activities: A Critical Review. *Sustainability*, 12(21), 9002. <https://doi.org/10.3390/su12219002>
- Brown, R. D., Vanos, J., Kenny, N., & Lenzholzer, S. (2015). Designing urban parks that ameliorate the effects of climate change. *Landscape and Urban Planning*, 138, 118–131. <https://doi.org/10.1016/j.landurbplan.2015.02.006>
- Brown-Amilian, S., & Akolade, Y. (2021). Disparities in COPD Hospitalizations: A Spatial Analysis of Proximity to Toxics Release Inventory Facilities in Illinois. *International Journal of Environmental Research and Public Health*, 18(24), 13128. <https://doi.org/10.3390/ijerph182413128>
- Brulle, R. J., & Pellow, D. N. (2006). Environmental Justice: Human Health and Environmental Inequalities. *Annual Review of Public Health*, 27(1), 103–124. <https://doi.org/10.1146/annurev.publhealth.27.021405.102124>
- Bulka, C., Nastoupil, L. J., Koff, J. L., Bernal-Mizrachi, L., Ward, K., Williams, J. N., Bayakly, A. R., Switchenko, J. M., Waller, L. A., & Flowers, C. R. (2016). Relations Between Residential Proximity to EPA-Designated Toxic Release Sites and Diffuse Large B-Cell Lymphoma Incidence. *Southern Medical Journal*, 109(10), 606–614. <https://doi.org/10.14423/SMJ.0000000000000545>
- Bullard, R. D. (1993). Anatomy of environmental racism and the environmental justice movement. *Confronting Environmental Racism: Voices from the Grassroots*, 15, 15–39.
- Bullard, R. D., Mohai, P., Saha, R., & Wright, B. (2008). Toxic Wastes and Race at Twenty: Why Race Still Matters after all of These Years. *Environmental Law*, 38, 371.
- Burgard, S. A., Seefeldt, K. S., & Zelner, S. (2012). Housing instability and health: Findings from the Michigan recession and recovery study. *Social Science & Medicine*, 75(12), 2215–2224. <https://doi.org/10.1016/j.socscimed.2012.08.020>
- Cakmak, S., Dales, R. E., & Vidal, C. B. (2007). Air Pollution and Mortality in Chile: Susceptibility among the Elderly. *Environmental Health Perspectives*, 115(4), 524–527. <https://doi.org/10.1289/ehp.9567>

California Office of Environmental Health Hazard Assessment. (2021). CalEnviroScreen 4.0 Report.

Canu, W. H., Jameson, J. P., Steele, E. H., & Denslow, M. (2017). Mountaintop Removal Coal Mining and Emergent Cases of Psychological Disorder in Kentucky. *Community Mental Health Journal*, 53(7), 802–810. <https://doi.org/10.1007/s10597-017-0122-y>

Centers for Disease Control and Prevention. (2020, August 21). Learn what could be triggering your asthma attacks. Centers for Disease Control and Prevention. <https://www.cdc.gov/asthma/triggers.html>

Chakraborty, J., Grineski, S. E., & Collins, T. W. (2019). Hurricane Harvey and people with disabilities: Disproportionate exposure to flooding in Houston, Texas. *Social Science & Medicine*, 226, 176–181. <https://doi.org/10.1016/j.socscimed.2019.02.039>

Chang, K. K. P., Wong, F. K. Y., Chan, K. L., Wong, F., Ho, H. C., Wong, M. S., Ho, Y. S., Yuen, J. W. M., Siu, J. Y., & Yang, L. (2020). The Impact of the Environment on the Quality of Life and the Mediating Effects of Sleep and Stress. *International Journal of Environmental Research and Public Health*, 17(22), 8529. <https://doi.org/10.3390/ijerph17228529>

Chen, X., & Kwan, M.-P. (2015). Contextual Uncertainties, Human Mobility, and Perceived Food Environment: The Uncertain Geographic Context Problem in Food Access Research. *American Journal of Public Health*, 105(9), 1734–1737. <https://doi.org/10.2105/AJPH.2015.302792>

Cohen, D. A., Marsh, T., Williamson, S., Derose, K. P., Martinez, H., Setodji, C., & McKenzie, T. L. (2010). Parks and physical activity: Why are some parks used more than others? *Preventive Medicine*, 50, S9–S12. <https://doi.org/10.1016/j.ypmed.2009.08.020>

Cohen, D. A., McKenzie, T. L., Sehgal, A., Williamson, S., Golinelli, D., & Lurie, N. (2007). Contribution of Public Parks to Physical Activity. *American Journal of Public Health*, 97(3), 509–514. <https://doi.org/10.2105/AJPH.2005.072447>

Coogan, P. F., White, L. F., Jerrett, M., Brook, R. D., Su, J. G., Seto, E., Burnett, R., Palmer, J. R., & Rosenberg, L. (2012). Air Pollution and Incidence of Hypertension and Diabetes Mellitus in Black Women Living in Los Angeles. *Circulation*, 125(6), 767–772. <https://doi.org/10.1161/CIRCULATIONAHA.111.052753>

Crouse, D. L., Peters, P. A., Hystad, P., Brook, J. R., van, D. A., Martin, R. V., Villeneuve, P. J., Jerrett, M., Goldberg, M. S., Pope, C. A., Brauer, M., Brook, R. D., Robichaud, A., Menard, R., & Burnett, R. T. (2015). Ambient PM_{2.5}, O₃, and NO₂ Exposures and Associations with Mortality over 16 Years of Follow-Up in the Canadian Census Health and Environment Cohort (CanCHEC). *Environmental Health Perspectives*, 123(11), 1180–1186. <https://doi.org/10.1289/ehp.1409276>

Cutter, S. L. (2012). *Hazards Vulnerability and Environmental Justice*. Routledge.

Cutter, S. L., Boruff, B. J., & Shirley, W. L. (2003). Social Vulnerability to Environmental Hazards*. *Social Science Quarterly*, 84(2), 242–261. <https://doi.org/10.1111/1540-6237.8402002>

Cutts, B. B., Darby, K. J., Boone, C. G., & Brewis, A. (2009). City structure, obesity, and environmental justice: An integrated analysis of physical and social barriers to walkable streets and park access. *Social Science & Medicine*, 69(9), 1314–1322. <https://doi.org/10.1016/j.socscimed.2009.08.020>

Dales, R. E., & Cakmak, S. (2016). Does Mental Health Status Influence Susceptibility to the Physiologic Effects of Air Pollution? A Population Based Study of Canadian Children. *PLOS ONE*, 11(12), e0168931. <https://doi.org/10.1371/journal.pone.0168931>

Deng, H., Eckel, S. P., Liu, L., Lurmann, F. W., Cockburn, M. G., & Gilliland, F. D. (2017). Particulate matter air pollution and liver cancer survival. *International Journal of Cancer*, 141(4), 744–749. <https://doi.org/10.1002/ijc.30779>

Dettenborn, L., Tietze, A., Bruckner, F., & Kirschbaum, C. (2010). Higher cortisol content in hair among long-term unemployed individuals compared to controls. *Psychoneuroendocrinology*, 35(9), 1404–1409. <https://doi.org/10.1016/j.psyneuen.2010.04.006>

Díaz, J., López-Bueno, J. A., López-Ossorio, J. J., González, J. L., Sánchez, F., & Linares, C. (2020). Short-term effects of traffic noise on suicides and emergency hospital admissions due to anxiety and depression in Madrid (Spain). *Science of The Total Environment*, 710, 136315. <https://doi.org/10.1016/j.scitotenv.2019.136315>

- Dockery, D. W., & Pope, C. A. (1994). Acute Respiratory Effects of Particulate Air Pollution. *Annual Review of Public Health*, 15(1), 107–132. <https://doi.org/10.1146/annurev.pu.15.050194.000543>
- Dong, G.-H., Qian, Z. (Min), Xaverius, P. K., Trevathan, E., Maalouf, S., Parker, J., Yang, L., Liu, M.-M., Wang, D., Ren, W.-H., Ma, W., Wang, J., Zelicoff, A., Fu, Q., & Simckes, M. (2013). Association Between Long-Term Air Pollution and Increased Blood Pressure and Hypertension in China. *Hypertension*, 61(3), 578–584. <https://doi.org/10.1161/HYPERTENSIONAHA.111.00003>
- Dórea, J. G. (2008). Persistent, bioaccumulative and toxic substances in fish: Human health considerations. *Science of The Total Environment*, 400(1), 93–114. <https://doi.org/10.1016/j.scitotenv.2008.06.017>
- Downey, L., & Van Willigen, M. (2005). Environmental Stressors: The Mental Health Impacts of Living Near Industrial Activity. *Journal of Health and Social Behavior*, 46(3), 289–305. <https://doi.org/10.1177/002214650504600306>
- Dubowsky, S. D., Suh, H., Schwartz, J., Coull, B. A., & Gold, D. R. (2006). Diabetes, Obesity, and Hypertension May Enhance Associations between Air Pollution and Markers of Systemic Inflammation. *Environmental Health Perspectives*, 114(7), 992–998. <https://doi.org/10.1289/ehp.8469>
- Ducatman, A., Ziemkiewicz, P., Quaranta, J., Vandivort, T., Mack, B., & VanAken, B. (2010). Coal slurry waste underground injection assessment. (Final report: Phase II. Prepared for the West Virginia Department of Health and Human Resources Office of Environmental Health Services. West Virginia University).
- Early, J., Davis, S. W., Quandt, S. A., Rao, P., Snively, B. M., & Arcury, T. A. (2006). Housing Characteristics of Farmworker Families in North Carolina. *Journal of Immigrant and Minority Health*, 8(2), 173–184. <https://doi.org/10.1007/s10903-006-8525-1>
- Elmenhorst, E.-M., Griefahn, B., Rolny, V., & Basner, M. (2019). Comparing the Effects of Road, Railway, and Aircraft Noise on Sleep: Exposure–Response Relationships from Pooled Data of Three Laboratory Studies. *International Journal of Environmental Research and Public Health*, 16(6), 1073. <https://doi.org/10.3390/ijerph16061073>
- Ernst, G. J. (1994). Racial and Economic Exploitation in the Siting of Toxic Wastes. *Bulletin of Science, Technology & Society*, 14(1), 28–32. <https://doi.org/10.1177/027046769401400105>
- Evans, G. W., & Kim, P. (2013). Childhood Poverty, Chronic Stress, Self-Regulation, and Coping. *Child Development Perspectives*, 7(1), 43–48. <https://doi.org/10.1111/cdep.12013>
- Fann, N., Lamson, A. D., Anenberg, S. C., Wesson, K., Risley, D., & Hubbell, B. J. (2012). Estimating the National Public Health Burden Associated with Exposure to Ambient PM_{2.5} and Ozone. *Risk Analysis*, 32(1), 81–95. <https://doi.org/10.1111/j.1539-6924.2011.01630.x>
- Faustman, E. M., Silbernagel, S. M., Fenske, R. A., Burbacher, T. M., & Ponce, R. A. (2000). Mechanisms underlying Children’s susceptibility to environmental toxicants. *Environmental Health Perspectives*, 108(suppl 1), 13–21. <https://doi.org/10.1289/ehp.00108s113>
- Foraster, M., Künzli N., Aguilera, I., Rivera, M., Agis, D., Vila, J., Bouso, L., Deltell, A., Marrugat, J., Ramos, R., Sunyer, J., Elosua, R., & Basagaña X. (2014). High Blood Pressure and Long-Term Exposure to Indoor Noise and Air Pollution from Road Traffic. *Environmental Health Perspectives*, 122(11), 1193–1200. <https://doi.org/10.1289/ehp.1307156>
- Forsyth, A. (2015). What is a walkable place? The walkability debate in urban design. *URBAN DESIGN International*, 20(4), 274–292. <https://doi.org/10.1057/udi.2015.22>
- Frank, L. D., Andresen, M. A., & Schmid, T. L. (2004). Obesity relationships with community design, physical activity, and time spent in cars. *American Journal of Preventive Medicine*, 27(2), 87–96. <https://doi.org/10.1016/j.amepre.2004.04.011>
- Frank, L. D., Sallis, J. F., Conway, T. L., Chapman, J. E., Saelens, B. E., & Bachman, W. (2006). Many Pathways from Land Use to Health: Associations between Neighborhood Walkability and Active Transportation, Body Mass Index, and Air Quality. *Journal of the American Planning Association*, 72(1), 75–87. <https://doi.org/10.1080/01944360608976725>
- Frank, L. D., Schmid, T. L., Sallis, J. F., Chapman, J., & Saelens, B. E. (2005). Linking objectively measured physical activity with objectively measured urban form: Findings from SMARTRAQ. *American Journal of Preventive Medicine*, 28(2, Supplement 2), 117–125. <https://doi.org/10.1016/j.amepre.2004.11.001>

- Frank, L., Kerr, J., Rosenberg, D., & King, A. (2010). Healthy Aging and Where You Live: Community Design Relationships With Physical Activity and Body Weight in Older Americans. *Journal of Physical Activity and Health*, 7(s1), S82–S90. <https://doi.org/10.1123/jpah.7.s1.s82>
- Freeman, L., Neckerman, K., Schwartz-Soicher, O., Quinn, J., Richards, C., Bader, M. D. M., Lovasi, G., Jack, D., Weiss, C., Konty, K., Arno, P., Viola, D., Kerker, B., & Rundle, A. G. (2013). Neighborhood walkability and active travel (walking and cycling) in New York City. *Journal of Urban Health: Bulletin of the New York Academy of Medicine*, 90(4), 575–585. <https://doi.org/10.1007/s11524-012-9758-7>
- Fuks, K. B., Weinmayr, G., Foraster, M., Dratva, J., Hampel, R., Houthuijs, D., Oftedal, B., Oudin, A., Panasevich, S., Penell, J., Sommar, J. N., Sørensen, M., Tiittanen, P., Wolf, K., Xun, W. W., Aguilera, I., Basagaña, X., Beelen, R., Bots, M. L., ... Hoffmann, B. (2014). Arterial Blood Pressure and Long-Term Exposure to Traffic-Related Air Pollution: An Analysis in the European Study of Cohorts for Air Pollution Effects (ESCAPE). *Environmental Health Perspectives*, 122(9), 896–905. <https://doi.org/10.1289/ehp.1307725>
- Gee, G. C., & Ponce, N. (2010). Associations Between Racial Discrimination, Limited English Proficiency, and Health-Related Quality of Life Among 6 Asian Ethnic Groups in California. *American Journal of Public Health*, 100(5), 888–895. <https://doi.org/10.2105/AJPH.2009.178012>
- Giorgini, P., Di Giosia, P., Grassi, D., Rubenfire, M., D. Brook, R., & Ferri, C. (2016). Air Pollution Exposure and Blood Pressure: An Updated Review of the Literature. *Current Pharmaceutical Design*, 22(1), 28–51.
- Go, A. S., Mozaffarian, D., Roger, V. L., Benjamin, E. J., Berry, J. D., Borden, W. B., Bravata, D. M., Dai, S., Ford, E. S., Fox, C. S., Franco, S., Fullerton, H. J., Gillespie, C., Hailpern, S. M., Heit, J. A., Howard, V. J., Huffman, M. D., Kissela, B. M., Kittner, S. J., ... Turner, M. B. (2013). Heart Disease and Stroke Statistics—2013 Update. *Circulation*, 127(1), e6–e245. <https://doi.org/10.1161/CIR.0b013e31828124ad>
- Goodling, E. (2020). Intersecting hazards, intersectional identities: A baseline Critical Environmental Justice analysis of US homelessness. *Environment and Planning E: Nature and Space*, 3(3), 833–856. <https://doi.org/10.1177/2514848619892433>
- Gray, S. C., Edwards, S. E., Schultz, B. D., & Miranda, M. L. (2014). Assessing the impact of race, social factors and air pollution on birth outcomes: A population-based study. *Environmental Health*, 13(1), 4. <https://doi.org/10.1186/1476-069X-13-4>
- Guarnieri, M., & Balmes, J. R. (2014). Outdoor air pollution and asthma. *The Lancet*, 383(9928), 1581–1592. [https://doi.org/10.1016/S0140-6736\(14\)60617-6](https://doi.org/10.1016/S0140-6736(14)60617-6)
- Haushofer, J., & Fehr, E. (2014). On the psychology of poverty. *Science*, 344(6186), 862–867. <https://doi.org/10.1126/science.1232491>
- Heikkala, E., Ala-Mursula, L., Taimela, S., Paananen, M., Vaaramo, E., Auvinen, J., & Karppinen, J. (2020). Accumulated unhealthy behaviors and psychosocial problems in adolescence are associated with labor market exclusion in early adulthood – a northern Finland birth cohort 1986 study. *BMC Public Health*, 20(1), 869. <https://doi.org/10.1186/s12889-020-08995-w>
- Helfand, G. E., & Peyton, L. J. (1999). A Conceptual Model of Environmental Justice. *Social Science Quarterly*, 80(1), 68–83.
- Hendryx, M., & Ahern, M. M. (2008). Relations Between Health Indicators and Residential Proximity to Coal Mining in West Virginia. *American Journal of Public Health*, 98(4), 669–671. <https://doi.org/10.2105/AJPH.2007.113472>
- Hendryx, M., Fedorko, E., & Anesetti-Rothermel, A. (2010). A geographical information system-based analysis of cancer mortality and population exposure to coal mining activities in West Virginia, United States of America. *Geospatial Health*, 4(2), 243. <https://doi.org/10.4081/gh.2010.204>
- Hendryx, M., & Luo, J. (2015). An examination of the effects of mountaintop removal coal mining on respiratory symptoms and COPD using propensity scores. *International Journal of Environmental Health Research*, 25(3), 265–276. <https://doi.org/10.1080/09603123.2014.938027>

- Hirsch, J. A., Moore, K. A., Barrientos-Gutierrez, T., Brines, S. J., Zagorski, M. A., Rodriguez, D. A., & Diez Roux, A. V. (2014). Built environment change and change in BMI and waist circumference: Multi-ethnic Study of Atherosclerosis. *Obesity, 22*(11), 2450–2457. <https://doi.org/10.1002/oby.20873>
- Hiscock, R., Macintyre, S., Ellaway, A., & Kearns, A. (2003). Residents and Residence: Factors Predicting the Health Disadvantage of Social Renters Compared to Owner-Occupiers. *Journal of Social Issues, 59*. <https://doi.org/10.1111/1540-4560.00076>
- Hong, Y.-C. (2013). Aging Society and Environmental Health Challenges. *Environmental Health Perspectives, 121*(3), a68–a69. <https://doi.org/10.1289/ehp.1206334>
- Houston, J. B., Hawthorne, J., Perreault, M. F., Park, E. H., Goldstein Hode, M., Halliwell, M. R., Turner McGowen, S. E., Davis, R., Vaid, S., McElderry, J. A., & Griffith, S. A. (2015). Social media and disasters: A functional framework for social media use in disaster planning, response, and research. *Disasters, 39*(1), 1–22. <https://doi.org/10.1111/disa.12092>
- Hynes, H. P., & Lopez, R. (2007). Cumulative Risk and a Call for Action in Environmental Justice Communities. *Journal of Health Disparities and Practice, 1*(2), 29–57.
- Jennings, V., Johnson Gaither, C., & Gragg, R. S. (2012). Promoting Environmental Justice Through Urban Green Space Access: A Synopsis. *Environmental Justice, 5*(1), 1–7. <https://doi.org/10.1089/env.2011.0007>
- Jerrett, M., Burnett, R. T., Beckerman, B. S., Turner, M. C., Krewski, D., Thurston, G., Martin, R. V., van Donkelaar, A., Hughes, E., Shi, Y., Gapstur, S. M., Thun, M. J., & Pope, C. A. (2013). Spatial Analysis of Air Pollution and Mortality in California. *American Journal of Respiratory and Critical Care Medicine, 188*(5), 593–599. <https://doi.org/10.1164/rccm.201303-0609OC>
- Jha, A., Lin, L., & Savoia, E. (2016). The Use of Social Media by State Health Departments in the US: Analyzing Health Communication Through Facebook. *Journal of Community Health, 41*(1), 174–179. <https://doi.org/10.1007/s10900-015-0083-4>
- Johnston, J., & MacDonald Gibson, J. (2015). Indoor Air Contamination from Hazardous Waste Sites: Improving the Evidence Base for Decision-Making. *International Journal of Environmental Research and Public Health, 12*(12), 15040–15057. <https://doi.org/10.3390/ijerph121214960>
- Kan, H., London, S. J., Chen, G., Zhang, Y., Song, G., Zhao, N., Jiang, L., & Chen, B. (2008). Season, Sex, Age, and Education as Modifiers of the Effects of Outdoor Air Pollution on Daily Mortality in Shanghai, China: The Public Health and Air Pollution in Asia (PAPA) Study. *Environmental Health Perspectives, 116*(9), 1183–1188. <https://doi.org/10.1289/ehp.10851>
- Khalid, N., Hussain, M., Young, H. S., Boyce, B., Aqeel, M., & Noman, A. (2018). Effects of road proximity on heavy metal concentrations in soils and common roadside plants in Southern California. *Environmental Science and Pollution Research, 25*(35), 35257–35265. <https://doi.org/10.1007/s11356-018-3218-1>
- Khan, A., Plana-Ripoll, O., Antonsen, S., Brandt, J., Geels, C., Landecker, H., Sullivan, P. F., Pedersen, C. B., & Rzhetsky, A. (2019). Environmental pollution is associated with increased risk of psychiatric disorders in the US and Denmark. *PLOS Biology, 17*(8), e3000353. <https://doi.org/10.1371/journal.pbio.3000353>
- Kiel, K., & Zabel, J. (2001). Estimating the Economic Benefits of Cleaning Up Superfund Sites: The Case of Woburn, Massachusetts. *The Journal of Real Estate Finance and Economics, 22*(2), 163–184. <https://doi.org/10.1023/A:1007835329254>
- Kim, D. Y., Staley, F., Curtis, G., & Buchanan, S. (2002). Relation Between Housing Age, Housing Value, and Childhood Blood Lead Levels in Children in Jefferson County, Ky. *American Journal of Public Health, 92*(5), 769–772. <https://doi.org/10.2105/AJPH.92.5.769>
- King, A. C., Sallis, J. F., Frank, L. D., Saelens, B. E., Cain, K., Conway, T. L., Chapman, J. E., Ahn, D. K., & Kerr, J. (2011). Aging in neighborhoods differing in walkability and income: Associations with physical activity and obesity in older adults. *Social Science & Medicine, 73*(10), 1525–1533. <https://doi.org/10.1016/j.socscimed.2011.08.032>

- Kouznetsova, M., Huang, X., Ma, J., Lessner, L., & Carpenter, D. O. (2007). Increased Rate of Hospitalization for Diabetes and Residential Proximity of Hazardous Waste Sites. *Environmental Health Perspectives*, 115(1), 75–79. <https://doi.org/10.1289/ehp.9223>
- Krivoshto, I. N., Richards, J. R., Albertson, T. E., & Derlet, R. W. (2008). The Toxicity of Diesel Exhaust: Implications for Primary Care. *The Journal of the American Board of Family Medicine*, 21(1), 55–62. <https://doi.org/10.3122/jabfm.2008.01.070139>
- Kwon, K., & Akar, G. (2022). People with disabilities and use of public transit: The role of neighborhood walkability. *Journal of Transport Geography*, 100, 103319. <https://doi.org/10.1016/j.jtrangeo.2022.103319>
- Lanphear, B. P., Weitzman, M., Winter, N. L., Eberly, S., Yakir, B., Tanner, M., Emond, M., & Matte, T. D. (1996). Lead-contaminated house dust and urban children's blood lead levels. *American Journal of Public Health*, 86(10), 1416–1421. <https://doi.org/10.2105/AJPH.86.10.1416>
- Last, J. A., Pinkerton, K. E., & Schelegle, E. S. (2017). Ozone and Oxidant Toxicity. In *Respiratory Toxicology* (Vols. 15–15, pp. 389–402). Elsevier Inc. <https://doi.org/10.1016/B978-0-08-100601-6.02076-7>
- Lee, C. (1992). Toxic Waste and Race in the United States. In *Race and the Incidence of Environmental Hazards*. Routledge.
- Lee, C. (2020). A Game Changer in the Making? Lessons from States Advancing Environmental Justice through Mapping and Cumulative Impact Strategies. *Environmental Law Reporter*, 50, 10203.
- Lin, H., Guo, Y., Zheng, Y., Zhao, X., Cao, Z., Rigdon, S. E., Xian, H., Li, X., Liu, T., Xiao, J., Zeng, W., Weaver, N. L., Qian, Z., Ma, W., & Wu, F. (2017). Exposure to ambient PM_{2.5} associated with overall and domain-specific disability among adults in six low- and middle-income countries. *Environment International*, 104, 69–75. <https://doi.org/10.1016/j.envint.2017.04.004>
- Lin, P., Lau, S. S. Y., Qin, H., & Gou, Z. (2017). Effects of urban planning indicators on urban heat island: A case study of pocket parks in high-rise high-density environment. *Landscape and Urban Planning*, 168, 48–60. <https://doi.org/10.1016/j.landurbplan.2017.09.024>
- Ljubimova, J. Y., Braubach, O., Patil, R., Chiechi, A., Tang, J., Galstyan, A., Shatalova, E. S., Kleinman, M. T., Black, K. L., & Holler, E. (2018). Coarse particulate matter (PM_{2.5-10}) in Los Angeles Basin air induces expression of inflammation and cancer biomarkers in rat brains. *Scientific Reports*, 8(1), 5708. <https://doi.org/10.1038/s41598-018-23885-3>
- Loh, M. M., Levy, J. I., Spengler, J. D., Houseman, E. A., & Bennett, D. H. (2007). Ranking Cancer Risks of Organic Hazardous Air Pollutants in the United States. *Environmental Health Perspectives*, 115(8), 1160–1168. <https://doi.org/10.1289/ehp.9884>
- Maantay, J. (2002). Zoning Law, Health, and Environmental Justice: What's the Connection? *Journal of Law, Medicine & Ethics*, 30(4), 572–593. <https://doi.org/10.1111/j.1748-720X.2002.tb00427.x>
- MacTavish, K., Eley, M., & Salamon, S. (2006). Housing Vulnerability among Rural Trailer-Park Households. *Georgetown Journal on Poverty Law and Policy*, 13, 95.
- Mahmood, A., O'Dea, E., Bigonnesse, C., Labbe, D., Mahal, T., Qureshi, M., & Mortenson, W. B. (2020). Stakeholders Walkability/Wheelability Audit in Neighbourhoods (SWAN): User-led audit and photographic documentation in Canada. *Disability & Society*, 35(6), 902–925. <https://doi.org/10.1080/09687599.2019.1649127>
- Maizlish, N., Delaney, T., Dowling, H., Chapman, D. A., Sabo, R., Woolf, S., Orndahl, C., Hill, L., & Snellings, L. (2019). California Healthy Places Index: Frames Matter. *Public Health Reports*, 134(4), 354–362. <https://doi.org/10.1177/0033354919849882>
- Malcoe, L. H., Lynch, R. A., Keger, M. C., & Skaggs, V. J. (2002). Lead sources, behaviors, and socioeconomic factors in relation to blood lead of native american and white children: A community-based assessment of a former mining area. *Environmental Health Perspectives*, 110(suppl 2), 221–231. <https://doi.org/10.1289/ehp.02110s2221>

- Marshall, E. G., Geary, N. S., Cayo, M. R., & Lauridsen, P. A. (1993). Residential exposure summary methodology for a reproductive health study of multiple hazardous waste sites. *Journal of Exposure Analysis and Environmental Epidemiology*, 3 Suppl 1, 87–98.
- Mawhorter, S., Crimmins, E. M., & Ailshire, J. A. (2021). Housing and cardiometabolic risk among older renters and homeowners. *Housing Studies*, 0(0), 1–23. <https://doi.org/10.1080/02673037.2021.1941792>
- McKee, A. M., & Cruz, M. A. (2021). Microbial and Viral Indicators of Pathogens and Human Health Risks from Recreational Exposure to Waters Impaired by Fecal Contamination. *Journal of Sustainable Water in the Built Environment*, 7(2), 03121001. <https://doi.org/10.1061/JSWBAY.0000936>
- Meltzer, R., & Schwartz, A. (2016). Housing Affordability and Health: Evidence From New York City. *Housing Policy Debate*, 26(1), 80–104. <https://doi.org/10.1080/10511482.2015.1020321>
- Meo, S. A., Memon, A. N., Sheikh, S. A., Rouq, F. A., Usmani, A. M., Hassan, A., & Arain, S. A. (n.d.). Effect of environmental air pollution on type 2 diabetes mellitus. *Diabetes Mellitus*, 6.
- Min, E., Gruen, D., Banerjee, D., Echeverria, T., Frelander, L., Schmeltz, M., Sagani, E., Piazza, M., Galaviz, V. E., Yost, M., & Seto, E. Y. W. (2019). The Washington State Environmental Health Disparities Map: Development of a Community-Responsive Cumulative Impacts Assessment Tool. *International Journal of Environmental Research and Public Health*, 16(22), 4470. <https://doi.org/10.3390/ijerph16224470>
- M. Nunes, L., Zhu, Y.-G., Y. Stigter, T., P. Monteiro, J., & R. Teixeira, M. (2011). Environmental impacts on soil and groundwater at airports: Origin, contaminants of concern and environmental risks. *Journal of Environmental Monitoring*, 13(11), 3026–3039. <https://doi.org/10.1039/C1EM10458F>
- Mohai, P., & Bryant, B. (1991). Race, Poverty & the Distribution of Environmental Hazards: Reviewing the Evidence. *Race, Poverty & the Environment*, 2(3/4), 3–27.
- Mohai, P., Lantz, P. M., Morenoff, J., House, J. S., & Mero, R. P. (2009). Racial and Socioeconomic Disparities in Residential Proximity to Polluting Industrial Facilities: Evidence From the Americans' Changing Lives Study. *American Journal of Public Health*, 99(S3), S649–S656. <https://doi.org/10.2105/AJPH.2007.131383>
- Mohai, P., & Saha, R. (2007). Racial Inequality in the Distribution of Hazardous Waste: A National-Level Reassessment. *Social Problems*, 54(3), 343–370. <https://doi.org/10.1525/sp.2007.54.3.343>
- Mohai, P., & Saha, R. (2015). Which came first, people or pollution? Assessing the disparate siting and post-siting demographic change hypotheses of environmental injustice. *Environmental Research Letters*, 10(11), 115008. <https://doi.org/10.1088/1748-9326/10/11/115008>
- Morello-Frosch, R., Zuk, M., Jerrett, M., Shamasunder, B., & Kyle, A. D. (2011). Understanding The Cumulative Impacts Of Inequalities In Environmental Health: Implications For Policy. *Health Affairs*, 30(5), 879–887. <https://doi.org/10.1377/hlthaff.2011.0153>
- Mulchandani, R., Smith, M., Armstrong, B., English National Study of Flooding and Health Study Group, Beck, C. R., & Oliver, I. (2019). Effect of Insurance-Related Factors on the Association between Flooding and Mental Health Outcomes. *International Journal of Environmental Research and Public Health*, 16(7), 1174. <https://doi.org/10.3390/ijerph16071174>
- Mullenbach, L. E., & Baker, B. L. (2020). Environmental Justice, Gentrification, and Leisure: A Systematic Review and Opportunities for the Future. *Leisure Sciences*, 42(5–6), 430–447. <https://doi.org/10.1080/01490400.2018.1458261>
- Münzel, T., Sørensen, M., & Daiber, A. (2021). Transportation noise pollution and cardiovascular disease. *Nature Reviews Cardiology*, 18(9), 619–636. <https://doi.org/10.1038/s41569-021-00532-5>
- Murgueytio, A. M., Evans, R. G., Sterling, D. A., Clardy, S. A., Shadel, B. N., & Clements, B. W. (1998). Relationship between Lead Mining and Blood Lead Levels in Children. *Archives of Environmental Health: An International Journal*, 53(6), 414–423. <https://doi.org/10.1080/00039899809605730>

- Neighbors, H. W., Caldwell, C., Williams, D. R., Nesse, R., Taylor, R. J., Bullard, K. M., Torres, M., & Jackson, J. S. (2007). Race, Ethnicity, and the Use of Services for Mental Disorders: Results From the National Survey of American Life. *Archives of General Psychiatry*, 64(4), 485–494. <https://doi.org/10.1001/archpsyc.64.4.485>
- Nepal, V., Banerjee, D., Perry, M., & Scott, D. (2012). Disaster Preparedness of Linguistically Isolated Populations: Practical Issues for Planners. *Health Promotion Practice*, 13(2), 265–271. <https://doi.org/10.1177/1524839910384932>
- Newman, S., & Holupka, C. S. (2016). Housing Affordability And Children's Cognitive Achievement. *Health Affairs*, 35(11), 2092–2099. <https://doi.org/10.1377/hlthaff.2016.0718>
- Nguyen, A., Mosadeghi, S., & Almario, C. V. (2017). Persistent digital divide in access to and use of the Internet as a resource for health information: Results from a California population-based study. *International Journal of Medical Informatics*, 103, 49–54. <https://doi.org/10.1016/j.ijmedinf.2017.04.008>
- Norris, G., YoungPong, S. N., Koenig, J. Q., Larson, T. V., Sheppard, L., & Stout, J. W. (1999). An association between fine particles and asthma emergency department visits for children in Seattle. *Environmental Health Perspectives*, 107(6), 489–493. <https://doi.org/10.1289/ehp.99107489>
- Ogneva-Himmelberger, Y., & Cooperman, B. (2010). Spatio-temporal Analysis of Noise Pollution near Boston Logan Airport: Who Carries the Cost? *Urban Studies*, 47(1), 169–182. <https://doi.org/10.1177/0042098009346863>
- Öhrström, E., Barregård, L., Andersson, E., Skånberg, A., Svensson, H., & Ångerheim, P. (2007). Annoyance due to single and combined sound exposure from railway and road traffic. *The Journal of the Acoustical Society of America*, 122(5), 2642–2652. <https://doi.org/10.1121/1.2785809>
- Ou, J. Y., Hanson, H. A., Ramsay, J. M., Kaddas, H. K., Pope, C. A., Leiser, C. L., VanDerslice, J., & Kirchhoff, A. C. (2020). Fine Particulate Matter Air Pollution and Mortality among Pediatric, Adolescent, and Young Adult Cancer Patients. *Cancer Epidemiology and Prevention Biomarkers*, 29(10), 1929–1939. <https://doi.org/10.1158/1055-9965.EPI-19-1363>
- Ozkurt, N., Hamamci, S. F., & Sari, D. (2015). Estimation of airport noise impacts on public health. A case study of zmir Adnan Menderes Airport. *Transportation Research Part D: Transport and Environment*, 36, 152–159. <https://doi.org/10.1016/j.trd.2015.02.002>
- Padula, A. M., Mortimer, K. M., Tager, I. B., Hammond, S. K., Lurmann, F. W., Yang, W., Stevenson, D. K., & Shaw, G. M. (2014). Traffic-related air pollution and risk of preterm birth in the San Joaquin Valley of California. *Annals of Epidemiology*, 24(12), 888–895.e4. <https://doi.org/10.1016/j.annepidem.2014.10.004>
- Pastor, M., Morello-Frosch, R., & Sadd, J. L. (2005). The Air Is Always Cleaner on the Other Side: Race, Space, and Ambient Air Toxics Exposures in California. *Journal of Urban Affairs*, 27(2), 127–148. <https://doi.org/10.1111/j.0735-2166.2005.00228.x>
- Pearsall, H. (2010). From Brown to Green? Assessing Social Vulnerability to Environmental Gentrification in New York City. *Environment and Planning C: Government and Policy*, 28(5), 872–886. <https://doi.org/10.1068/c08126>
- Peek, L., & Stough, L. M. (2010). Children With Disabilities in the Context of Disaster: A Social Vulnerability Perspective. *Child Development*, 81(4), 1260–1270. <https://doi.org/10.1111/j.1467-8624.2010.01466.x>
- Peel, J. L., Tolbert, P. E., Klein, M., Metzger, K. B., Flanders, W. D., Todd, K., Mulholland, J. A., Ryan, P. B., & Frumkin, H. (2005). Ambient Air Pollution and Respiratory Emergency Department Visits. *Epidemiology*, 16(2), 164–174.
- Pellow, D. N. (2021). Struggles for Environmental Justice in US Prisons and Jails. *Antipode*, 53(1), 56–73. <https://doi.org/10.1111/anti.12569>
- Perkins, H. A., Heynen, N., & Wilson, J. (2004). Inequitable access to urban reforestation: The impact of urban political economy on housing tenure and urban forests. *Cities*, 21(4), 291–299. <https://doi.org/10.1016/j.cities.2004.04.002>
- Peters, A., Dockery, D. W., Muller, J. E., & Mittleman, M. A. (2001). Increased Particulate Air Pollution and the Triggering of Myocardial Infarction. *Circulation*, 103(23), 2810–2815. <https://doi.org/10.1161/01.CIR.103.23.2810>

- Petri, D., Licitra, G., Vigotti, M. A., & Fredianelli, L. (2021). Effects of Exposure to Road, Railway, Airport and Recreational Noise on Blood Pressure and Hypertension. *International Journal of Environmental Research and Public Health*, 18(17), 9145. <https://doi.org/10.3390/ijerph18179145>
- Phillips, L. A., Solís, P., Wang, C., Varfalameyeva, K., & Burnett, J. (2021). Engaged Convergence Research: An Exploratory Approach to Heat Resilience in Mobile Homes. *The Professional Geographer*, 73(4), 619–631. <https://doi.org/10.1080/00330124.2021.1924805>
- Pierce, G., & Jimenez, S. (2015). Unreliable Water Access in U.S. Mobile Homes: Evidence From the American Housing Survey. *Housing Policy Debate*, 25(4), 739–753. <https://doi.org/10.1080/10511482.2014.999815>
- Pope III, C. A., Burnett, R. T., Thun, M. J., Calle, E. E., Krewski, D., Ito, K., & Thurston, G. D. (2002). Lung Cancer, Cardiopulmonary Mortality, and Long-term Exposure to Fine Particulate Air Pollution. *JAMA*, 287(9), 1132–1141. <https://doi.org/10.1001/jama.287.9.1132>
- Rajagopalan, S., Al, -Kindi Sadeer G., & Brook, R. D. (2018). Air Pollution and Cardiovascular Disease. *Journal of the American College of Cardiology*, 72(17), 2054–2070. <https://doi.org/10.1016/j.jacc.2018.07.099>
- Reuben, A., Rutherford, G. W., James, J., & Razani, N. (2020). Association of neighborhood parks with child health in the United States. *Preventive Medicine*, 141, 106265. <https://doi.org/10.1016/j.ypmed.2020.106265>
- Reynolds, P., Von, B. J., Gunier, R. B., Goldberg, D. E., Hertz, A., & Smith, D. F. (2003). Childhood cancer incidence rates and hazardous air pollutants in California: An exploratory analysis. *Environmental Health Perspectives*, 111(4), 663–668. <https://doi.org/10.1289/ehp.5986>
- Rigolon, A. (2017). Parks and young people: An environmental justice study of park proximity, acreage, and quality in Denver, Colorado. *Landscape and Urban Planning*, 165, 73–83. <https://doi.org/10.1016/j.landurbplan.2017.05.007>
- Rigolon, A., Browning, M., & Jennings, V. (2018). Inequities in the quality of urban park systems: An environmental justice investigation of cities in the United States. *Landscape and Urban Planning*, 178, 156–169. <https://doi.org/10.1016/j.landurbplan.2018.05.026>
- Ristovski, Z. D., Miljevic, B., Surawski, N. C., Morawska, L., Fong, K. M., Goh, F., & Yang, I. A. (2012). Respiratory health effects of diesel particulate matter. *Respirology*, 17(2), 201–212. <https://doi.org/10.1111/j.1440-1843.2011.02109.x>
- Roemmich, J. N., Johnson, L., Oberg, G., Beeler, J. E., & Ufholz, K. E. (2018). Youth and Adult Visitation and Physical Activity Intensity at Rural and Urban Parks. *International Journal of Environmental Research and Public Health*, 15(8), 1760. <https://doi.org/10.3390/ijerph15081760>
- Sadler, R. C., LaChance, J., & Hanna-Attisha, M. (2017). Social and Built Environmental Correlates of Predicted Blood Lead Levels in the Flint Water Crisis. *American Journal of Public Health*, 107(5), 763–769. <https://doi.org/10.2105/AJPH.2017.303692>
- Schirnding, Y. von, Mathee, A., Kibel, M., Robertson, P., Strauss, N., & Blignaut, R. (2003). A study of pediatric blood lead levels in a lead mining area in South Africa. *Environmental Research*, 93(3), 259–263. [https://doi.org/10.1016/S0013-9351\(03\)00117-8](https://doi.org/10.1016/S0013-9351(03)00117-8)
- Schultz, B. D., Morara, M., Buxton, B. E., & Weintraub, M. (2017). Predicting Blood-Lead Levels Among U.S. Children at the Census Tract Level. *Environmental Justice*, 10(5), 129–136. <https://doi.org/10.1089/env.2017.0005>
- Sergeev, A. V., & Carpenter, D. O. (2005). Hospitalization Rates for Coronary Heart Disease in Relation to Residence Near Areas Contaminated with Persistent Organic Pollutants and Other Pollutants. *Environmental Health Perspectives*, 113(6), 756–761. <https://doi.org/10.1289/ehp.7595>
- Shapiro, M. D. (2005). Equity and information: Information regulation, environmental justice, and risks from toxic chemicals. *Journal of Policy Analysis and Management*, 24(2), 373–398. <https://doi.org/10.1002/pam.20094>
- Shcherbatykh, I., Huang, X., Lessner, L., & Carpenter, D. O. (2005). Hazardous waste sites and stroke in New York State. *Environmental Health*, 4(1), 18. <https://doi.org/10.1186/1476-069X-4-18>

- Shishegar, N. (2014). The impact of green areas on mitigating urban heat island effect: A review. *International Journal of Environmental Sustainability*, 9(1), 119–130. <https://doi.org/10.18848/2325-1077/CGP/v09i01/55081>
- Shrader-Frechette, K. (2002). *Environmental Justice: Creating Equality, Reclaiming Democracy*. Oxford University Press.
- Shrestha, R., Flacke, J., Martinez, J., & Van Maarseveen, M. (2016). Environmental Health Related Socio-Spatial Inequalities: Identifying “Hotspots” of Environmental Burdens and Social Vulnerability. *International Journal of Environmental Research and Public Health*, 13(7), 691. <https://doi.org/10.3390/ijerph13070691>
- Slaughter, J. C., Kim, E., Sheppard, L., Sullivan, J. H., Larson, T. V., & Claiborn, C. (2005). Association between particulate matter and emergency room visits, hospital admissions and mortality in Spokane, Washington. *Journal of Exposure Science & Environmental Epidemiology*, 15(2), 153–159. <https://doi.org/10.1038/sj.jea.7500382>
- Smedley, B. D. (2012). The Lived Experience of Race and Its Health Consequences. *American Journal of Public Health*, 102(5), 933–935. <https://doi.org/10.2105/AJPH.2011.300643>
- Smith, L. (2015). Walkability Audit Tool. *Workplace Health & Safety*, 63(9), 420–420. <https://doi.org/10.1177/2165079915595307>
- Stieb, D. M., Burnett, R. T., Beveridge, R. C., & Brook, J. R. (1996). Association between ozone and asthma emergency department visits in Saint John, New Brunswick, Canada. *Environmental Health Perspectives*, 104(12), 1354–1360. <https://doi.org/10.1289/ehp.961041354>
- Stowe, E. W., Hughey, S. M., Hallum, S. H., & Kaczynski, A. T. (2019). Associations between Walkability and Youth Obesity: Differences by Urbanicity. *Childhood Obesity*, 15(8), 555–559. <https://doi.org/10.1089/chi.2019.0063>
- Sturm, R., & Cohen, D. (2014). Proximity to Urban Parks and Mental Health. *The Journal of Mental Health Policy and Economics*, 17(1), 19–24.
- Suglia, S. F., Duarte, C. S., & Sandel, M. T. (2011). Housing Quality, Housing Instability, and Maternal Mental Health. *Journal of Urban Health*, 88(6), 1105–1116. <https://doi.org/10.1007/s11524-011-9587-0>
- Sutherland, R. A., & Tolosa, C. A. (2001). Variation in Total and Extractable Elements with Distance from Roads in an Urban Watershed, Honolulu, Hawaii. *Water, Air, and Soil Pollution*, 127(1), 315–338. <https://doi.org/10.1023/A:1005283932003>
- Tanzer, R., Malings, C., Hauryliuk, A., Subramanian, R., & Presto, A. A. (2019). Demonstration of a Low-Cost Multi-Pollutant Network to Quantify Intra-Urban Spatial Variations in Air Pollutant Source Impacts and to Evaluate Environmental Justice. *International Journal of Environmental Research and Public Health*, 16(14), 2523. <https://doi.org/10.3390/ijerph16142523>
- Tate, A., & Emrich, C. T. (2021). Assessing social equity in disasters. *Eos*, 102.
- Teron, L. (2016). Sustainably Speaking: Considering Linguistic Isolation in Citywide Sustainability Planning. *Sustainability: The Journal of Record*, 9(6), 289–294. <https://doi.org/10.1089/sus.2016.29072.it>
- Thayamballi, N., Habiba, S., Laribi, O., & Ebisu, K. (2021). Impact of Maternal Demographic and Socioeconomic Factors on the Association Between Particulate Matter and Adverse Birth Outcomes: A Systematic Review and Meta-analysis. *Journal of Racial and Ethnic Health Disparities*, 8(3), 743–755. <https://doi.org/10.1007/s40615-020-00835-2>
- Tolbert, J., Nov 06, A. D. P., & 2020. (2020, November 6). Key Facts about the Uninsured Population. KFF. <https://www.kff.org/uninsured/issue-brief/key-facts-about-the-uninsured-population/>
- Tolbert, P. E., Mulholland, J. A., Macintosh, D. L., Xu, F., Daniels, D., Devine, O. J., Carlin, B. P., Klein, M., Butler, A. J., Nordenberg, D. F., Frumkin, H., Ryan, P. B., & White, M. C. (2000). Air Quality and Pediatric Emergency Room Visits for Asthma and Atlanta, Georgia. *American Journal of Epidemiology*, 151(8), 798–810. <https://doi.org/10.1093/oxfordjournals.aje.a010280>
- United States Environmental Protection Agency. (2019). *EJSCREEN Technical Documentation*.
- US Environmental Protection Agency. (2014). *Environmental Quality Index: Overview Report*.

- US Environmental Protection Agency. (2016). Regulatory Impact Analysis Accidental Release Prevention Requirements: Risk Management Programs Under the Clean Air Act, Section 112(r)(7).
- US Environmental Protection Agency. (2017). National Walkability Index: Methodology and User Guide.
- US Environmental Protection Agency. (2021). Public Comment on EPA's National Enforcement Initiatives for Fiscal Years 2017–2019. 80 FR 5535.
- Villeneuve, P. J., Chen, L., Rowe, B. H., & Coates, F. (2007). Outdoor air pollution and emergency department visits for asthma among children and adults: A case-crossover study in northern Alberta, Canada. *Environmental Health*, 6(1), 40. <https://doi.org/10.1186/1476-069X-6-40>
- Voelkel, J., Hellman, D., Sakuma, R., & Shandas, V. (2018). Assessing Vulnerability to Urban Heat: A Study of Disproportionate Heat Exposure and Access to Refuge by Socio-Demographic Status in Portland, Oregon. *International Journal of Environmental Research and Public Health*, 15(4), 640. <https://doi.org/10.3390/ijerph15040640>
- Weuve, J., Kaufman, J. D., Szpiro, A. A., Curl, C., Puett, R. C., Beck, T., Evans, D. A., & Mendes, de L. C. F. (2016). Exposure to Traffic-Related Air Pollution in Relation to Progression in Physical Disability among Older Adults. *Environmental Health Perspectives*, 124(7), 1000–1008. <https://doi.org/10.1289/ehp.1510089>
- White-Newsome, J., O'Neill, M. S., Gronlund, C., Sunbury, T. M., Brines, S. J., Parker, E., Brown, D. G., Rood, R. B., & Rivera, Z. (2009). Climate Change, Heat Waves, and Environmental Justice: Advancing Knowledge and Action. *Environmental Justice*, 2(4), 197–205. <https://doi.org/10.1089/env.2009.0032>
- Whitworth, K. W., Symanski, E., & Coker, A. L. (2008). Childhood Lymphohematopoietic Cancer Incidence and Hazardous Air Pollutants in Southeast Texas, 1995–2004. *Environmental Health Perspectives*, 116(11), 1576–1580. <https://doi.org/10.1289/ehp.11593>
- Wichmann, H.-E. (2007). Diesel Exhaust Particles. *Inhalation Toxicology*, 19(sup1), 241–244. <https://doi.org/10.1080/08958370701498075>
- Williams, D. R., & Mohammed, S. A. (2009). Discrimination and racial disparities in health: Evidence and needed research. *Journal of Behavioral Medicine*, 32(1), 20–47. <https://doi.org/10.1007/s10865-008-9185-0>
- Wilson, S. M., Fraser-Rahim, H., Williams, E., Zhang, H., Rice, L., Svendsen, E., & Abara, W. (2012). Assessment of the Distribution of Toxic Release Inventory Facilities in Metropolitan Charleston: An Environmental Justice Case Study. *American Journal of Public Health*, 102(10), 1974–1980. <https://doi.org/10.2105/AJPH.2012.300700>
- Wolch, J. R., Byrne, J., & Newell, J. P. (2014). Urban green space, public health, and environmental justice: The challenge of making cities 'just green enough.' *Landscape and Urban Planning*, 125, 234–244. <https://doi.org/10.1016/j.landurbplan.2014.01.017>
- Wong, R., Harris, J. K., Staub, M., & Bernhardt, J. M. (2017). Local Health Departments Tweeting About Ebola: Characteristics and Messaging. *Journal of Public Health Management and Practice*, 23(2), e16–e24. <https://doi.org/10.1097/PHH.0000000000000342>
- Wood, L., Hooper, P., Foster, S., & Bull, F. (2017). Public green spaces and positive mental health – investigating the relationship between access, quantity and types of parks and mental wellbeing. *Health & Place*, 48, 63–71. <https://doi.org/10.1016/j.healthplace.2017.09.002>
- Woolhandler, S., & Himmelstein, D. U. (2017). The Relationship of Health Insurance and Mortality: Is Lack of Insurance Deadly? *Annals of Internal Medicine*, 167(6), 424–431. <https://doi.org/10.7326/M17-1403>
- Wright, R. J., Rodriguez, M., & Cohen, S. (1998). Review of psychosocial stress and asthma: An integrated biopsychosocial approach. *Thorax*, 53(12), 1066–1074. <https://doi.org/10.1136/thx.53.12.1066>
- Wu, C., Wu, S., Wu, Y.-H., Cullen, A. C., Larson, T. V., Williamson, J., & Liu, L.-J. S. (2009). Cancer risk assessment of selected hazardous air pollutants in Seattle. *Environment International*, 35(3), 516–522. <https://doi.org/10.1016/j.envint.2008.09.009>
- Yi, O., Kim, H., & Ha, E. (2010). Does area level socioeconomic status modify the effects of PM10 on preterm delivery? *Environmental Research*, 110(1), 55–61. <https://doi.org/10.1016/j.envres.2009.10.004>